

doi: 10.15407/ujpe60.08.0795

B. ESHCHANOV, SH. OTAJONOV, N. SOLIEVA, A. ISAMATOV

Department of Physics, National University of Uzbekistan
(Tashkent 100174, Uzbekistan; e-mail: bakhodir.eshchanov@gmail.com)

INTENSITY DISTRIBUTION IN THE SPECTRUM OF MOLECULAR LIGHT SCATTERING AND RELAXATION EFFECTS IN LIQUIDS

PACS 77.22.Ej, 78.35.+c

The structure and shape of the spectral bands of orientational molecular scattering of light in toluene are analyzed. The possibility of their use for obtaining information about the features of interactions of molecules is revealed. It is shown that the observed scattering regularities can be explained, by basing on the mechanism of hindered rotation of the molecules with regard for the specific intermolecular interaction in toluene.

Keywords: orientation spectroscopy, molecular scattering of light, orientation motion of molecules in liquids, relaxation mechanism, relaxation time, fluctuation.

1. Introduction

The problem of determining the nature and strength of intermolecular interactions in condensed media, as well as the most common regularities of changes in the optical spectra of molecular complexes, due to the influence of intermolecular interactions of different nature is the fundamental branch of modern condensed matter spectroscopy [1–7].

An intermolecular spectrum can be considered as a source of information on the specificity of the condensed state of matter, since it reflects the motion of molecules, intermolecular interactions, and the temperature variation of the structure of matter. Analysis of the experimental works shows that, with rare exceptions, they do not address the full range of the intermolecular spectrum over a wide temperature range. This work is devoted to the study of the temperature behavior of the depolarized component of the molecular scattering of light (DCMSL) over a wide temperature interval 20–200 °C and a frequency interval from 0 to 50–60 cm⁻¹ in toluene. For this purpose, in accordance with certain assumptions, it is required to observe the thermal behavior of a number of parameters of the condensed state of matter; to analyze the possibility of obtaining information on the nature of the temperature change in the dynamics of movement and interaction of molecules of

matter and its structural changes on the temperature change of the DCMSL spectrum.

2. The Experimental Part

The DCMSL spectra were recorded on the experimental setup constructed on the base of a DFS-12 spectrometer, which is used to obtain the spectra of weak signals operating in the photon counting mode. The description of the installation can be found in [8]. The substances used, depending on the grade of purity, were cleaned, dried, de-dusted, and outgassed by double vacuum distillation. The purity of the object was controlled by the relative intensities of the shifted and unshifted Mandelstam–Brillion components of the scattering polarized part.

3. Results and Discussion

The object of studies was toluene (C₆H₅CH₃) – asymmetric on the molecular polarizability. Data on the DCMSL spectrum of toluene are very limited. In particular, there are no data on the temperature dependence of the light scattering, on the shape of the spectrum, etc. The intensity distribution of high-frequency areas (50–200 cm⁻¹) had previously been studied in [8]. As a result of the statistical processing of the distant areas of the recorded DCMSL contour by various analytical functions, it is shown that the best approximation corresponds to a Gaussian. Therefore, it was interesting to apply the dynamics of temperature changes for the low-frequency




Fig. 1. Reciprocal spectral density of DCMSL lines in toluene molecules as a function of the squared frequency: at a temperature of 20 °C (1), at a temperature of 200 °C (2)

region of the DCMSL spectrum of toluene to establish the mechanism of scattering in the frequency interval 0–50 cm^{-1} .

Experiments show that the observed line contour in DCMSL (in the coordinates $1/\tilde{\gamma}-\nu^2$, where $\tilde{\gamma}$ is the spectral density of the contour) represents the sum of two dispersive lines in the frequency interval 0–60 cm^{-1} , with the interface between the components equal to about 15 cm^{-1} at the temperature $T = 20^\circ\text{C}$. With the heating, the dispersive line undergoes a further simplification, and the boundary of the band shifts toward lower frequencies, and this point is about 7 cm^{-1} at $T = 200^\circ\text{C}$ (Fig. 1).

Toluene is structurally related to an asymmetric top. According to the model of hindered rotation of molecules (MHRM) [5,9], the DCMSL contour is a superposition of three dispersion contours, whose half-widths (σ_i) are determined by three relaxation times τ_i , $\sigma_i = \frac{1}{2\pi c\tau_i}$, and the weight (α_i) is defined via the main value of the components for the molecule polarizability tensor

$$\alpha_i = \frac{(a_k - a_j)^2}{2\gamma^2}, \quad (1)$$

where $\gamma^2 = \frac{1}{2}[(a_1 - a_2)^2 + (a_1 - a_3)^2 + (a_2 - a_3)^2]$ is the anisotropy of molecules, a_1, a_2, a_3 – the principal values of components of the polarizability tensor of a molecule.

According to MHRM, a contour in the liquids with non-axisymmetric molecules must consist of three Lorentzians. However, the structure of a toluene molecule allows suggesting that a Lorentz half-width σ_2 related to the rotation about the axis passing through the substituted vertices of the benzene ring a_1 (Fig. 2) should be much more than the half-widths of two other components. Furthermore, the theoretical value of contour weight α_3 associated with the rotation about the third axis (the axis of symmetry for toluene) calculated in the “gas” approximation is much smaller than α_2 .

Because of this expansion, the DCMSL contour consisting of two, rather than three Lorentz components can be considered a reasonable approximation. Subsequently, the first contour means a sum of the first and third contours.

Table 1 shows the weights α_1^{theor} , α_2^{theor} , and α_3^{theor} calculated by formula (1). Data on the main values of polarizabilities are taken from [10]. As we can see from this table, the weight of the third contour α_3 is very small (4%), so the observed DCMSL contour can be represented as the sum of

Table 1

Substance	α_1^{theor}	α_2^{theor}	α_3^{theor}
$\text{C}_6\text{H}_5\text{CH}_3$	0.61	0.35	0.04




Fig. 2. Structural formula and the choice of axes of a molecule of toluene

the two dispersion (Lorentzian) contours (“narrow” – in the frequency interval $0\text{--}15\text{ cm}^{-1}$ and “wide” – $15\text{--}50\text{ cm}^{-1}$).

Table 2 shows the results of the processing of the experimental DCMSL contours obtained by the decomposition of the experimental contour into components by the method of successive approximations.

As we can see from this table, σ_1 , σ_2 , α_1 , α_2 are parameters of the contour components, \mathfrak{J} is the integrated intensity of the total contour in relative units. For σ_2 and α_2 , the accuracy estimation was performed by the method of least squares. For σ_1 and \mathfrak{J} , the relative mean square errors of the mean for all the measurement cycle are given.

Analysis of Table 2 shows that the narrow contour σ_1 indicates the strong temperature dependence – $\frac{\sigma_1(200^\circ\text{C})}{\sigma_1(20^\circ\text{C})} = 3.83$; the wide contour σ_2 weakly depends on the temperature – $\frac{\sigma_2(200^\circ\text{C})}{\sigma_2(20^\circ\text{C})} = 1.24$.

Comparison of the theoretical and experimental values of weight for contour components shows that, at room temperature, the values of weights α_1 and

α_2 are different. With increasing the temperature, the difference is reduced; the further increase of the temperature leads to a divergence again. This may be due to the fact that, in the process of scattering, not only the rotational motion is observed, but also other types of motion, in particular, the vibrational one.

With increasing the temperature, the half-widths of contours increase, and a narrower contour σ_1 expands relatively faster, than a wide contour. The total integrated intensity of DCMSL increases with the temperature. The most interesting result is that the intensities $\mathfrak{J}_1 = \mathfrak{J} \alpha_1$ and $\mathfrak{J}_2 = \mathfrak{J} \alpha_2$, of the narrow and broad components of the Lorentz contours have different signs in the temperature range: at the beginning, they increase with the temperature, and then they decrease. In view of Anselm’s theory [11], this corresponds to a positive correlation of molecules on one axis and a negative one on the other one, which results in significant structure details of the short-range order in the liquid.

The temperature dependence of the intensity of these components will be determined by changes in the correlation of the orientations along the corresponding axes. Thus, the temperature measurements allow, in principle, obtaining enough information about the nature of the short-range order in liquids. As a result, a task of direct evaluation of the role of correlations of the intensities of individual components for the anisotropic scattering becomes very urgent. Without such a calculation, the conclusions about the correlations obtained based on [11] from the temperature dependence of the scattering have a conditional character.

Table 2

$T, ^\circ\text{C}$	σ_1, cm^{-1}	σ_2, cm^{-1}	α_1	α_2	\mathfrak{J}
20	1.20	10.00	0.67	0.33	8000
50	2.00	11.00	0.62	0.37	8500
100	2.60	12.50	0.60	0.40	8800
150	4.60	12.40	0.54	0.46	9000
200	4.60	12.40	0.51	0.49	9300
The statistical error	± 0.1	± 0.5	± 0.02	± 0.02	1-1.5

- V.E. Pogorelov, L.A. Bulavin, I. Doroshenko, O. Fesjun, and O. Veretennikov, *J. of Mol. Struct.* **708**, No. 1-3, 61 (2014).
- N.A. Atamas, A.M. Yaremko, L.A. Bulavin, V.E. Pogorelov, S. Berski, Z. Latajka, H. Ratajczak, and A. Abkowitz-Bieńko, *J. of Mol. Struct.* **605**, 187 (2002).
- I.I. Adamenko, L.A. Bulavin, V. Ilyin, S.A. Zelinsky, and K.O. Moroz, *J. of Mol. Liquids* **127**, 90 (2006).
- L.A. Bulavin, A.I. Fisenko, and N.P. Malomuzh, *Chem. Phys. Lett.* **453**, 183 (2008).
- P.K. Khabibullaev, L.A. Bulavin, V.E. Pogorelov, F.X. Tuhvatullin, A.I. Lizengevich, Sh.O. Otajonov, and A.J. Jumabaev, *The Dynamics of Molecules of the Liquid* (Fan, Tashkent, 2009).

6. I.Yu. Doroshenko, V.Ye. Pogorelov, G.A. Pitsevich, and V. Shablinskas, *The Dynamics of Molecules of the Liquid* (LAMBERT Academic Publ., 2012).
7. L.A. Bulavin, V.Ya. Gotsulskiy, and V.E. Chechko, *Ukr. J. Phys.* **59**, 881 (2014).
8. Sh. Otajonov, B. Eshchanov, and A. Isamatov, *Ukr. J. Phys.* **59**, 254 (2014).
9. Sh. Otajonov, *Doctoral thesis* (Tashkent, 1992).
10. I.L. Fabelinskii, *Molecular Scattering of Light* (Nauka, Moscow, 1965) (in Russian).
11. A.I. Anselm, *JETP* **17**, 439 (1947).

Received 01.05.15

Б.Х. Ешчанов, Ш. Отажонов, Н. Солиева, А. Исаматов

РОЗПОДІЛ ІНТЕНСИВНОСТІ В СПЕКТРІ
МОЛЕКУЛЯРНОГО РОЗСІЮВАННЯ СВІТЛА
І РЕЛАКСАЦІЙНІ ЯВИЩА В РІДИНАХ

Резюме

Проаналізовано структуру та форму смуг орієнтаційних спектрів молекулярного розсіювання світла в толуолі. Виявлено можливості їх використання для отримання інформації про особливості взаємодії молекул. Показано, що закономірності, які спостерігаються в розсіянні, можна пояснити, виходячи з механізму загальмованого обертання молекул рідини з урахуванням специфіки міжмолекулярної взаємодії в толуолі.